



# Preliminary results for sensitivity analyses of CH4 emissions derived from ground-based synthetic observations through the TM5 4D-VAR system

Maria Gabriella Villani<sup>1</sup>
Peter Bergamaschi<sup>1</sup>
Jan Fokke Meirink<sup>2</sup>
Maarten Krol<sup>2,3</sup>

- [1] European Commission Joint Research Centre, Institute for Environment and Sustainability, I-21020 Ispra (Va), Italy
- [2] IMAU, Utrecht, The Netherlands
- [3] Wageningen University and Research Centre, Wageningen, the Netherlands





## **Experiment set-up**



- Use of 4D-Var system
- global spatial resolution (6° by 4°)
- year 2003
- 11 emission categories


# **Create Synthetic data** ("Observations")

-"continuous measurements (3hr)"

#### **Perturbations on emissions**


- emissions from rice cultivation: 50% less of the true values

#### STATIONS LOCATION










-4.00 -6.00 ₺



























Category contribution: (Emis<sub>cat\_spos</sub>-Emis<sub>cat\_true</sub>)/ (Emis<sub>tot\_spos</sub>-Emis<sub>tot\_true</sub>)





Category contribution: (Emis<sub>cat\_spos</sub>-Emis<sub>cat\_true</sub>)/ (Emis<sub>tot\_spos</sub>-Emis<sub>tot\_true</sub>)



### **Conclusions**



 4D-VAR system can retrieve the CH4 "true emissions" using ground-based observations

Issues to consider for this experiment:

- Emissions attributed to categories emitting at the same location
- Emissions attributed to categories having different temporal scale correlations


Work in progress and future analyses:

- Flask measurements
- Extending the study to other emission categories



#### **WORK IN PROGRESS**






"Continuous Measurements"


"Flask Measurements"










































12